

Numerical results for the Anderson transition. Comment

I. M. Suslov

P.L.Kapitza Institute for Physical problems

117334, Moscow, Russia

E-mail: suslov@kapitza.ras.ru

Abstract

Answer to cond-mat/0106005, cond-mat/0106006 and additional notes are given concerning my previous comment (cond-mat/0105325).

Recently I have discussed the numerical results for the Anderson transition [1]. Objections [2] and [3] have appeared after that. We answer here to the arguments of [2, 3] and give the additional comments.

1. I did not state in [1] that $\nu = 1$ (in fact, my most probable value was 1.25). I tried to show that a problem exists: numerical results contradict to all other information on the critical exponents. It is not my personal problem, and it is not my personal affair to solve it. Indeed, a value $\nu \approx 1$ is desirable from viewpoint of this other information and I tried to understand, is not it possible to agree it with the raw numerical data by the change of interpretation and by more realistic estimation of errors. I found it possible for the data that I considered as the best. May be I was a little wrong in the latter estimation.

Authors of [2] argue that I used the systems of too small size (28^3), "while currently sizes $\sim 50^3$ (for ELS) and $18^2 \times 10^8$ (for TM) are standard"¹. A size $18^2 \times 10^8$ should kick a man out his feet, while authors of [2] know perfectly well that only the minimal length scale ($L = 18$) is relevant. In fact, such systems *are small*.² I do not know about "currently" and "standard" but only one result is cited in [2] for systems $\sim 50^3$ [6]. With my procedure used, authors of [2] obtained this result with uncertainty $\nu = 1.25 \div 1.75$, which is not drastically different from uncertainty $0.8 \div 1.7$ that I have derived from Zharekeshev and Kramer data [7]. So, it is admitted that the largest scale data give uncertainty $1.25 \div 1.75$ and not 1% [5]. At the present stage, I am quite satisfied with such progress.

¹ Abbreviations of [2]: TM and ELS — transfer matrix and energy level statistics methods, FSS — finite size scaling.

² I can understand such trick in the text, where authors of [2] are in the course of discussion, but the same thing is present in the figure capture: "the system sizes of TM data are larger than for ELS data".

2. Authors of [2] claim, that my procedure is not interesting because it does not give increase of accuracy. They are mistaken here: I had not purpose to increase accuracy. I wanted to dig out the real uncertainty of results. According to [2], estimation of error given by my procedure approximately corresponds to a scattering of results due to different authors. So this estimation is reliable and reasonably conservative. I can hardly dream of this. By the way, only "people with experience in FSS" can believe in accuracy of several percents, looking at Fig. 1 of [2].

In fact, authors of [2] have overlooked the essence of my suggestions. I have introduced some function $f(L)$, in order to have a convenient language for discussion of scaling corrections. My suggestions are simple. Show your function $f(L)$, so that we can have a look at it. Then it will be evident, has this function so good power-law behavior as you say, or not. It will be clear, is there change of the effective ν with the length scale, or not. Then we can compare such functions for different quantities and different models, in order to estimate a systematic error. Then we can see, how this error change with L and so on.

I do not insist, that my procedure for determination of the function $f(L)$ is the best. In fact, it is useful in the case of rather poor raw data. If these data are sufficiently detailed and accurate, it is more preferable to extract the function $f(L)$ by direct linearization (in fact, it was admitted in [1]). Such procedure was used in a number of papers and I have no priority in it. But it was only recently, that explicit plots of such functions have appeared [8, 9].

3. I agree with [3], that in the general case a fixed point depends on L and irrelevant parameters are uncritical. The fixed point μ^* has a regular dependence in a/L and the main correction is the ordinary surface effect. With this dependence taken into account, my Eq.(5) takes a form

$$Q(\mu) = F \{ \mu^*(L/a), A_1(\tau)s^{y_1}, A_2(\tau)s^{y_2}, A_3(\tau)s^{y_3}, \dots \} , \quad (5')$$

Putting $A_1(\tau) = b_1\tau$, $A_i(\tau) = c_i + b_i\tau$ ($i \geq 2$), we have instead (6)

$$Q(\tau, L) = F \{ a/L, b_1\tau(L/a)^{y_1}, (c_2 + b_2\tau)(L/a)^{y_2}, (c_3 + b_3\tau)(L/a)^{y_3}, \dots \} , \quad (6')$$

For $L \gg a$ we have instead (7) and (8) correspondingly

$$\begin{aligned} Q(\tau, L) &= [F \{ 0, 0, 0, 0, \dots \} + B_0(a/L) + B_2(L/a)^{y_2} + B_3(L/a)^{y_3}, \dots] + \\ &+ \tau [C_1(L/a)^{y_1} + C_2(L/a)^{y_2} + C_3(L/a)^{y_3}, \dots] \equiv F_0(L/a) + \tau f(L/a) , \quad \tau(L/a)^{y_1} \ll 1 \end{aligned} \quad (7')$$

and

$$Q(\tau, L) \approx F \{ 0, b_1\tau(L/a)^{y_1}, 0, 0, \dots \} \equiv G \{ \tau(L/a)^{y_1} \} , \quad \tau(L/a)^{y_1} \gtrsim 1 . \quad (8')$$

Correction to the critical point

$$\tilde{\tau} = \tau + \Delta(a/L) , \quad \Delta(a/L) = [F_0(L/a) - F_0(0)] / f(L/a)$$

is essential for small τ in Eq.(7') but is irrelevant in Eq.(8'). So unification (7') and (8') has a form:

$$Q(\tau, L) = G \{ \tilde{\tau} f(L/a) \} . \quad (9')$$

One should shift the curves for different L , so as they have a common intersection point, and use the same procedure as in [1]. Such shift is routinely used in the most of papers. There was no need in it for the data of [7].

For $d = 4 - \epsilon$ the first irrelevant exponent y_2 is of the order of ϵ [10] and linearization in the parameter $c_2(L/a)^{y_2}$ is invalid for $L \lesssim L_0 \sim a \exp\{const/\epsilon\}$. The corresponding dependence on L is inessential in the restricted intervals of L but there is a slow drift of results on the large scale L_0 . May be, it is a clue to the whole problem.

4. There are a lot of vague speculations in [3] concerning a number of nonlinear parameters. Authors were able to calculate that in my procedure this number is even greater than in their one. In spite of these speculations, it is evident to anyone that

(a) all steps of my procedure are well defined and unambiguous.

(b) authors of [3] did not give a clear answer to a clear question: how do they avoid ambiguity of their procedure?

With modifications of [3], my procedure becomes ambiguous too. There is no sense to discuss the results of the unclear procedure.

5. One can carry out a very simple experiment. Take some function, f.e. $f(x) = x^2/(1+x)$, add some noise to it, and try to find its asymptotic behavior Ax^α at $x \rightarrow \infty$, using only values of $f(x)$ for $x = 1, 2, \dots, 20$. You will find it rather difficult to formulate a procedure that gives the reliable and reasonably conservative estimation of errors. The more or less successful procedure is as follows. Take an interval $(x_{min}, 20)$ and increase x_{min} until a simplest fit $\log f(x) = C + \alpha \log x$ give for χ^2 its normal value $\approx n$, where n is a number of points³. It is a necessary condition for a systematic error to be of the order of a statistical one. Now increase x_{min} *a little more*: then the formal statistical error will give more or less reasonable estimation of the real error. *A little more* is rather subjective and depends on the degree of conservatism. This procedure is successful, if a noise is uncorrelated and you have reliable estimation of its amplitude. In the general case such procedure is minimal and gives only the lower bound of the error. In my experience, related with a paper [11] and its development [12], the noise was strongly correlated and a formal statistical error was typically two orders less than a real one (even for the good values of χ^2). I can only wonder that some people are so confident in their accuracy.

The given example is simple, because corrections to asymptotics *are regular*. You can increase accuracy of α by many orders, fitting $\log f(x)$ as $C_0 + \alpha \log x + C_1/x + C_2/x^2 + \dots$ (this fit is linear and, consequently, unique). It is not the case that is interesting for us.

If a structure of corrections is power-like but unknown, you should use a nonlinear fit $f(x) = Ax^\alpha + A_1x^{\alpha_1} + A_2x^{\alpha_2} + \dots$ and you immediately meet with a problem of many minima. Their origin is rather simple. Take an arbitrary succession $\alpha, \alpha_1, \alpha_2 \dots$ (such

³ We suppose that this number is much greater than a number of parameters.

that $\alpha > \alpha_1 > \alpha_2 > \dots$) and use for $f(x)$ a linear fit $Ax^\alpha + A_1x^{\alpha_1} + A_2x^{\alpha_2} + \dots$ (only $A, A_1, A_2, \dots$ are changed). It will be successful with sufficient number of terms and χ^2 takes its normal value. Take this result as "a zero approximation" and use the general nonlinear fit (not only $A, A_1, A_2, \dots$ but also $\alpha, \alpha_1, \alpha_2, \dots$ are changed). The quantity χ^2 cannot be essentially lower than its normal value, and can change only slightly. So, there is a local minimum for it, close to "a zero approximation". Taking different successions $\alpha, \alpha_1, \alpha_2, \dots$, you can generate a great number of such minima. A value of α is different in different successions. So you can obtain an *arbitrary* result for it⁴ by such procedure.

I think, it is clear now, why one should not believe in any results obtained with such fits. If a structure of corrections is unknown, one hardly can invent something better than the simplest procedure given in beginning of this section.

6. After [1] I had a constructive discussion with P. Markoš⁵ and now I have better understanding of a general situation. Here are my impressions of it.

(i) There is a good paper [9], where the transfer-matrix method and the system sizes up to $L = 24$ were used. The raw data are rather detailed and accurate, so a function $f(L)$ can be obtained with good accuracy⁶ by direct linearization of W -dependence. More than that, Fig. 2 of this paper shows a number of such functions (in the log-log coordinates) for the different Lyapunov exponents. Some observations can be made in this figure. For $L \lesssim 10$ the slopes of the curves differ very strongly and correspond to values of ν in the interval $0.8 \div 1.5$. For $L \gtrsim 10$ practically all curves change their slopes and the more or less unique slope (with $\nu \approx 1.5$) arises. Nevertheless, the curve for the least exponent z_1 (which, according to [9], should reach its asymptotics most quickly) has a tendency to change its slope in the opposite direction. For $L > 12$, this slope is compatible with a value $\nu = 1.25$, if a point for $L = 24$ (evident outlier) is excluded. One can see, that a possible change of the effective ν with L is not a fantasy.

(ii) There is a lot of high precision results for $L \lesssim 15$ and they indeed give a value $\nu \approx 1.5$. Let us assume, that this value is not true, but effective. Then it should change with increase of L . In practice, one can double [7, 9] or triple [6] this size, though there are not many such attempts. With such increase of L , a situation cannot improve drastically. One can be able only to see some change of ν with L . There is a number of reasons, why this change could not be observed reliably:

(a) Accuracy of the raw data is worse by an order of magnitude for $L \gtrsim 15$.

(b) A need for a careful treatment was underestimated. The researches were satisfied that the large scale data roughly correspond to well-established results for small L , though the most of fits are not satisfactory.

(c) When linearization of W -dependence was used, the interval of linearization was kept fixed. It gives the wrong tendency due to increase of nonlinear effects for the large L (see

⁴ It is restricted only by the assumptions made.

⁵ He has no responsibility for my conclusions.

⁶ I think, this accuracy is a little exaggerated. One can see in Fig. 2 of [9] that a number of points deviates from the general smooth dependence in a quantity that is not controlled by the given error.

footnote 5 in [1]). This tendency is not large in magnitude, but the whole situation is rather unstable. One can see in Fig. 2 of [9] that a shift of the large L data in one standard deviation produces essential change of results.

As a consequence, the large L data appeared to be *practically useless*. In fact, this conclusion is supported by authors of [2], who refer primarily to results for small and not for large L .

(iii) If the raw data have a quality comparable with [9], the adequate procedure should contain, in our opinion, the following stages:

(a) In order to have the efficient use of the χ^2 procedure, the statistical uncertainties of the raw data are not simply estimated, but are calculated independently for each point in the course of some formal procedure. Probably, it can be done with the use of different realizations of a random potential.

(b) The function $f(L)$ is obtained by linearization of W -dependence. The interval of linearization is adjusted in accordance with χ^2 and the error is properly estimated. It should be done independently for each L , but with the use of the same routine. More conservatism is desirable at this stage.

(c) The function $f(L)$ is plotted. It is fitted by the power law dependence in the interval (L_{min}, L_{max}) where L_{min} is adjusted with the use of χ^2 and L_{max} is the largest L that is possible to reach. Minimization of χ^2 is a clever procedure and it provides a necessary compromise between the high accuracy data for small L and the poor accuracy data for large L : the latter are properly weighted, but all useful information is extracted of them. Adjustment of L_{min} allows to exclude the main body of a systematic error, which is contained in the small L data. Excess of conservatism at the stage (b) is partially compensated, because smaller L_{min} becomes admissible.

(d) The rest of a systematic error can be controlled by comparison of results for different quantities and different models.

I have impression, that nobody had used this procedure in a full extent⁷.

(iv) Some comments concerning the 4D case. The most advanced results are those of Zharekeshev and Kramer [8]. With $\nu = 1$, they have a parameter $\tau(L/a)^{1/\nu}$ (in designations of [1]) of the order unity, and a condition for linearization of W -dependence is marginally fulfilled. With $\nu = 1/2$ (as it should be), the above parameter is ~ 10 and they are deeply in the nonlinear regime. So, a possibility $\nu = 1/2$, in fact, was not tested numerically.

Some confirmation of these arguments can be found in [14]. Fig. 1,a of this paper shows W -dependences for another quantity (Lyapunov exponent, but not level statistics) which are essentially more detailed in comparison with [8]. Indeed, they are strikingly nonlinear. More attention should be given to investigation of these nonlinearities.

This work was financially supported by INTAS (Grant 99-1070) and RFBR (00-02-17129).

⁷ Sensitivity of results to restriction of the intervals in W and L was observed by MacKinnon[13].

References

- [1] I. M. Suslov, cond-mat/0105325
- [2] P. Cain, M. L. Ndawana, R. A. Römer, M. Schreiber, cond-mat/0106005
- [3] K. Slevin, T. Ohtsuki, cond-mat/0106006
- [4] M. Schreiber, H. Grussbach, Phys. Rev. Lett. **76**, 1687 (1996).
- [5] K. Slevin, T. Ohtsuki, Phys. Rev. Lett. **82**, 382 (1999).
- [6] F. Milde, R. A. Römer, M. Schreiber, cond-mat/9909210, cond-mat/0009469.
- [7] I. Kh. Zharekeshev, B. Kramer, Phys. Rev. Lett. **79**, 717 (1997).
- [8] I. Kh. Zharekeshev, B. Kramer, Ann. Phys. (Leipzig) **7**, 442 (1998).
- [9] P. Markoš, cond-mat/9907413.
- [10] I. M. Suslov, JETP **86**, 798 (1998), cond-mat/0007027
- [11] I. M. Suslov, JETP Lett. **71**, 217 (2000), hep-ph/0006066.
- [12] I. M. Suslov, JETP **93**, N 7 (2001), in print.
- [13] A. MacKinnon, J. Phys.: Condens. Matter **6**, 2511 (1994).
- [14] P. Markoš, M. Heneke, J. Phys.: Condens. Matter **6**, L765 (1994).